

ABOUT FOLLICULAR LYMPHOMA

JUST THE FACTS

Lymphoma is a cancer of the lymphatic system, which is a system of lymph nodes found throughout the body, including the neck, underarms, chest, abdomen and groin/pelvis.¹


The lymphatic system is part of the body's immune system, transporting white blood cells (T-cells and B-cells) to **fight infection in the body.**¹



In some instances, the B-cells can be locked in a state of replication, **which can become malignant over time.**³

Of the estimated 74,000 individuals diagnosed with NHL in the United States each year, FL accounts for approximately

20% of cases, or about 14,800 individuals.⁴



One form of non-Hodgkin lymphoma (NHL) is **follicular lymphoma (FL)**,² **which originates in B-cells** (which make antibodies to fight infection).¹

WHO IS DIAGNOSED WITH FL?



According to the National Cancer Institute, **FL is most common among white men and women**, compared with other ethnic groups.⁵



24.7% of patients have Stage 1 disease at diagnosis, meaning the disease is located in one area of the body.⁵



27% of patients present with Stage 4 disease, meaning the disease has spread throughout the body.⁵

55-64

is the age range of people most frequently diagnosed with FL, with a median age of 63.⁵

SYMPTOMS & DIAGNOSIS

Many patients are asymptomatic at diagnosis,⁶ but some common symptoms they may experience include⁷:



Painless, swollen lymph nodes in the neck, armpit or groin



Persistent fatigue, abdominal or chest pain

Fever, night sweats or unexplained weight loss



A **diagnosis of FL is made following a biopsy of a patient's lymph node tissue** and may also include additional tests, including a CT scan to determine if the disease has spread to other parts of the body.

LIVING WITH FL

- The highest five-year relative survival rate occurs in patients 55 years of age and those diagnosed with Stage 1 disease.⁵ The **median overall survival is 13-14 years from diagnosis.**^{8,9}
- While patients with FL may live for many years, **FL remains a typically incurable disease**; patients will often experience periods of remission before their disease returns (relapse).
- Patients may **also become resistant to previous treatments** they have received (refractory) as their disease becomes more aggressive and/or difficult to treat.
- Given many will undergo multiple rounds of therapy; patients should work closely with their health care provider to **develop long-term treatment plans** that take into account severity of disease and lifestyle needs.



REFERENCES

1. Lymphoma - Non-Hodgkin: Introduction. Cancer.Net. Accessed July 6, 2020. Available at: <https://www.cancer.net/cancer-types/lymphoma-non-hodgkin/introduction>
2. Lymphoma - Non-Hodgkin: Subtypes. Cancer.Net. Accessed July 6, 2020. Available at: <https://www.cancer.net/cancer-types/lymphoma-non-hodgkin/subtypes>
3. Béguelin W, et al. EZH2 Is Required for Germinal Center Formation and Somatic EZH2 Mutations Promote Lymphoid Transformation. Cancer Cell Article. 2013;23(5): 677-692. doi: 10.1016/j.ccr.2013.04.011.
4. About Non-Hodgkin Lymphoma. American Cancer Society. Accessed July 6, 2020. Available at: <https://www.cancer.org/cancer/non-hodgkin-lymphoma/about.html>
5. Cancer Stat Facts: NHL - Follicular Lymphoma. National Cancer Institute, SEER. Available at: <https://seer.cancer.gov/statfacts/html/follicular.html>

6. Understanding Follicular Lymphoma: Relapsed/Refractory. Lymphoma Research Foundation. Accessed July 27, 2021. https://lymphoma.org/wp-content/uploads/2020/09/LRF_Fact-sheet_Follicular-Lymphoma-Relapsed_Refractory_090920.pdf
7. Lymphoma. Mayo Clinic. Accessed July 6, 2020. Available at: <https://www.mayoclinic.org/diseases-conditions/lymphoma/symptoms-causes/syc-20352638>
8. Tan D, et al. Improvements in observed and relative survival in follicular grade 1-2 lymphoma during 4 decades: the Stanford University experience. Blood. 2013;122(6):981-987. doi: 10.1182/blood-2013-03-491514.
9. Liu Q, et al. Improvement of Overall and Failure-Free Survival in Stage IV Follicular Lymphoma: 25 years of treatment experience at The University of Texas M.D. Anderson Cancer Center. J Clin Oncol. 2006;24(10):1582-9. doi: 10.1200/JCO.2005.03.3696.